

Helsinki 11.2.2004

RECEIVED

01 MAR 2004

WIPO PCT

ETUOIKEUSTODISTUS
PRIORITY DOCUMENT



Hakija
Applicant

Oy Jurilab Ltd
Kuopio

Patenttihakemus nro
Patent application no

20022178

Tekemispäivä
Filing date

11.12.2002

Kansainvälinen luokka
International class

C12Q

Keksinnön nimitys
Title of invention

"Method and kit for detecting a risk of diabetes"
(Menetelmä ja testipakkaus diabetesriskin havaitsemiseksi)

Täten todistetaan, että oheiset asiakirjat ovat tarkkoja jäljennöksiä Patentti- ja rekisterihallitukselle alkuaan annetuista selityksestä, patenttivaatimuksista ja tiivistelmästä.

This is to certify that the annexed documents are true copies of the description, claims and abstract originally filed with the Finnish Patent Office.

Marketta Tehikoski

Marketta Tehikoski
Apulaistarkastaja

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Maksu 50 €
Fee 50 EUR

Maksu perustuu kauppa- ja teollisuusministeriön antamaan asetukseen 1027/2001 Patentti- ja rekisterihallituksen maksullisista suoritteista muutoksineen.

The fee is based on the Decree with amendments of the Ministry of Trade and Industry No. 1027/2001 concerning the chargeable services of the National Board of Patents and Registration of Finland.

Osoite: Arkadiankatu 6 A Puhelin: 09 6939 500
P.O.Box 1160 Telephone: + 358 9 6939 500
FIN-00101 Helsinki, FINLAND

Telefax: 09 6939 5328
Telefax: + 358 9 6939 5328

Method and kit for detecting a risk of diabetes

FIELD OF THE INVENTION

5 The invention provides a method and kit for detecting or diagnosing a risk of or predisposition to type 2 diabetes or a metabolic syndrome in a subject, the method comprising the steps of providing a biological sample of the subject to be tested and detecting the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in the biological sample. The invention also relates to a method for the treatment of
10 type 2 diabetes.

BACKGROUND OF THE INVENTION

There are two sub-types of α -adrenoceptors: the α_1 -adrenoceptors (α_1 -ARs), on the effector organs (postsynaptic) and the α_2 -adrenoceptors (α_2 -ARs) on the nerve endings (presynaptic). The α_2 -adrenoceptors mediate many of the physiological effects of
15 epinephrine and norepinephrine including a reduction of release of norepinephrine. Three genetic subtypes of α_2 -ARs are known in humans, denoted as α_{2A} -, α_{2B} - and α_{2C} -AR that are located on chromosomes 10, 2 and 4, respectively (Calzada and Artinano 2001). The tissue distributions and physiological and pharmacological functions of the receptor subtypes have been reviewed elsewhere (Docherty 1998, Calzada and Artinano
20 2001). Based on recent studies with gene-targeted mice, α_{2A} -ARs mediate most of the pharmacological actions ascribed to currently available α_2 -AR agonists, including inhibition of neurotransmitter release, central hypotensive and bradycardic effects, sedation and anaesthesia, and analgesia (MacMillan et al 1998).

The pancreatic islets are richly innervated by parasympathetic, sympathetic and sensory
25 nerves. Several different neurotransmitters are stored within the terminals of these nerves, the classical neurotransmitters, acetylcholine and norepinephrine, and neuropeptides, which exert local effects. Stimulation of the autonomic nerves affects islet hormone secretion. Thus, insulin secretion is stimulated by parasympathetic nerves or their neurotransmitters and inhibited by sympathetic nerves or their
30 neurotransmitters. The islet autonomic nerves seem to be of physiological importance

allowing oscillations of islet hormone secretion and are also involved in the islet adaptation in disease states (Ahrén 2000).

Sympathetic nervous system (SNS) over-activity plays a key role in the development of the metabolic syndrome, which is characterised by the combination of high blood pressure, increased glucose production, decreased glucose utilisation, increase in triglycerides and VLDL, decrease in HDL and insulin resistance (Borchard 2001). Insulin resistance is defined as an impaired biologic response to exogenous or endogenous insulin and is a state in which the ability of insulin to suppress hepatic glucose output and stimulate the uptake and utilisation of glucose by muscle and adipose tissues is impaired (American Diabetes Association 1998).

The effects of the SNS on glucose and lipid metabolism are mediated by circulating catecholamines (epinephrine and norepinephrine) and the direct sympathetic innervation of the liver, adipose tissues and skeletal muscle. It has been generally recognised that increased sympathetic neural activity (SNA) produces catabolic effects on glucose and lipid metabolism (Nonogaki 2000) and inhibits insulin release from pancreatic beta-cells and glucose uptake into skeletal muscle cells, which are a major site of insulin-mediated glucose uptake (Borchard 2001) and a primary site of insulin resistance as measured by euglycaemic glucose clamp technique (Moan et al. 1996). Increased SNA causes vasoconstriction of the skeletal muscle arteries leading to impaired glucose tolerance and to insulin resistance.

In isolated adipocytes beta-adrenergic stimulation induces a rapid down-regulation of insulin receptors together with a decrease in insulin-mediated glucose transport. Insulin resistance leads to a breakdown of stored triglycerides in the adipose tissue and an increase in plasma free fatty acids. As a consequence, hepatic synthesis of triglycerides from free fatty acids and conversion of triglycerides to VLDL-cholesterol is enhanced. Catecholamines may further increase lipolysis in adipocytes which results in an elevated release of free fatty acids into the blood stream. Free fatty acids decrease glucose-stimulated insulin release from the pancreas, which further enhances glucose intolerance. Furthermore, catecholamines may inhibit lipoprotein lipase and thus increase VLDL, which is linked to a decrease in HDL (Borchard 2001). The end result

is a steady state of hyperinsulinemia. Support for this concept is based on these observations: a) a decreased skeletal muscle capillary density has been found in insulin-resistant states of hypertension, obesity, and type 2 diabetes; b) antihypertensive drugs that cause vasoconstriction worsen insulin resistance and those that cause vasodilation improve insulin sensitivity; and c) exercise training improves insulin sensitivity and increases skeletal muscle capillary density (Julius et al. 1991).

The role of SNS in insulin resistance and diabetes is further strengthened by studies in which blockade of parasympathetic activity has been shown to increase the prevalence and severity of diabetes in mice suggesting role of unopposed sympathetic activity (Kvist-Reimer et al. 2002) and sympathectomy abrogated the development of both hyperinsulinemia and hypertension in fructose hypertensive rats - a widely used model to study the inter-relationship between hyperinsulinemia, insulin resistance and high blood pressure (Verma et al. 1999).

In addition to the findings that increased SNA promotes insulin resistance and type 2 diabetes there is also experimental evidence that hyperinsulinemia may itself increase sympathetic neural outflow by activating anteroventral portion of the third ventricle in the central nervous system, a region implicated in arterial pressure regulation and sympathetic neural control. In rats and humans acute insulin elicits increased lumbar SNA along with elevated plasma norepinephrine and studies have firmly established in both humans and experimental animals that insulin administration produces marked increases in SNA directed largely to skeletal muscle tissues (Muntzel 1999). Insulin is also involved in the initiation of appropriate changes in sympathetic outflow within the central nervous system in response to diet. During fasting state, the small decline in the plasma glucose concentration and the larger decrease in the plasma insulin concentration result in diminished insulin-mediated uptake and metabolism of glucose to carbon dioxide by insulin-sensitive cells in the ventromedial hypothalamus. The decreased glucose metabolism stimulates the activity of an inhibitory pathway between the hypothalamus and the brain stem, suppressing tonically active sympathetic centres in the brain stem and decreasing central sympathetic activity. Conversely, in insulin-resistant states, the increase in plasma glucose and the greater increase in plasma insulin (hyperinsulinemia) stimulate insulin-mediated uptake and metabolism of glucose by the hypothalamic cells.

The increase in glucose metabolism diminishes the activity of the inhibitory pathway, disinhibiting tonically active brain-stem centres and increasing central sympathetic activity leading to hypertension by stimulating the heart, the vasculature, and the kidneys (Reaven et al. 1996).

- 5 Thus it can be concluded that increased SNA and hyperinsulinemia by a mutually dependent process results in insulin resistance and increased risk of type 2 diabetes and carries a high mortality and risk of cardiovascular diseases, kidney failure and infectious complications as measured by resistance and pulsatility indices (Takahashi et al. 1998; Valensi et al. 1997; Weinrauch et al. 1995; Weston et al. 1999) and warrants
10 an early detection and treatment.

- Both β -adrenoceptor antagonists (β -blockers) and diuretics have been observed to influence glucose balance unfavourably, elevating blood glucose and serum insulin levels. Drug-induced hyperglycemia is a growing concern. Several antihypertensive drugs have an adverse effect on glucose tolerance that may partially or completely
15 negate the beneficial effects of reducing blood pressure as it relates to the incidence of coronary heart disease and its complications. Diuretics and beta-blockers have the greatest adverse effect on glucose intolerance (Houston 1986) and patients receiving beta-blocker treatment or diuretics may be at increased risk for developing hyperglycemia and subsequent diabetes mellitus (Luna et al. 2001; Bengtsson 1984;
20 Gress et al. 2000; Bengtsson et al. 1984).

- Metabolic studies of beta-blockers and diuretics aroused concern about the diabetogenic potential of these drugs. Subsequently, the results of prospective cohort studies and clinical trials suggested a causal link between the use of beta-blockers or diuretics and
25 the subsequent development of type 2 diabetes (Gress et al. 2000; Bengtsson et al. 1984; Lithell 1998; Papaccio et al. 1987). The risk of diabetes was 28 percent greater among those who took beta-blockers than among those who took no medication, without regard to the presence or absence of hypertension, sociodemographic characteristics, health-related behaviour, family history with respect to diabetes, and a
30 variety of coexisting conditions (Gress et al. 2000). Likewise, some observational studies have identified higher estimates of relative risk: for instance, in one study,

subjects who took beta-blockers had up to 6.1 times the risk of diabetes of those who did not (Samuelsson et al. 1994).

5 Thiazide diuretic-induced hyperglycemia occurs primarily through the reduction in total body potassium and the subsequent decreased insulin secretion (Luna et al. 2001; Greenberg 2000; Perez-Stable et al. 1983) by the beta cells, and reductions in extracellular fluid volume and cardiac output. This is compounded by increases in catecholamines from sympathetic nerve activity, which decrease peripheral glucose utilisation (Wilcox 1999). Potential mechanisms by which beta-blockers may contribute
10 to the development of diabetes include weight gain, attenuation of the beta-receptor-mediated release of insulin from pancreatic beta cells, and decreased blood flow through the microcirculation in skeletal-muscle tissue, leading to decreased insulin sensitivity (Sowers et al. 2000).

15 SUMMARY OF THE INVENTION

The object of this invention is to provide a screening method to assess if an individual is at risk to develop diabetes or a metabolic syndrome based on the genotype of α_{2B} -adrenoceptor (ADRA2B) gene. The invention also provides a method for the treatment
20 of type 2 diabetes in a human or animal subject. A further object of the invention is a method to determine whether a subject will benefit of different antihypertensive treatments with regard to their effects on the glucose balance and metabolism. Another object of the invention is a method for the selection of subjects for clinical trials testing antidiabetogenic drugs and compounds with effects on the insulin sensitivity.

25 The present invention concerns a method for detecting a risk of diabetes or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor i.e. to determine if said subject's genotype of the human α_{2B} -adrenoceptor is variant type, comprising the steps of

- a) providing a biological sample of the subject to be tested,

- b) detecting the presence or absence of variant genotype of the human α_{2B} -adrenoceptor gene in the biological sample, the presence of variant genotype indicating an increased risk of diabetes in said subject.

5 According to the invention, the method allows the determination whether said subject is of said variant genotype or not, a presence of said variant genotype in the biological sample, such as a blood sample or a buccal swab sample, thus indicating an increased risk of the subject to develop diabetes, and/or indicating the subject is in need for treatment, such as α_{2B} -selective or α_{2B} -nonselective antagonist therapy.

10 Preferably, the present invention concerns a method for detecting a risk of diabetes or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant α_2 -adrenoceptor gene i.e. to determine if said subject's genotype of the α_2 -adrenoceptor gene is of the deletion/deletion (D/D) type, comprising the steps of

- 15 a) providing a biological sample of the subject to be tested,
b) detecting the presence of α_2 -adrenoceptor deletion/deletion (D/D) type in the biological sample, the presence of D/D genotype indicating an increased risk of diabetes or a metabolic syndrome in said subject or the subject's need for α_{2B} -selective or α_{2B} -nonselective antagonist therapy for
20 diabetes.

A further object of the invention is a method for treating diabetes in a diabetic or glucose intolerant subject by reducing the sympathetic tone or activity, lowering blood or tissue norepinephrine or epinephrine concentrations and/or antagonizing α_{2B} -
25 adrenoceptors of the human or animal subject.

The present invention is also directed to a kit for detecting a risk of type 2 diabetes or a metabolic syndrome in a subject, comprising means for determining the pattern of alleles encoding a variant α_{2B} -adrenoceptor in a biological sample.

In the spirit of the invention, it is also conceivable that any variation in the α_{2B} -adrenoceptor gene that alters the structure or function of the mature α_{2B} -adrenoceptor protein can be used in predicting the risk of type 2 diabetes or a metabolic syndrome, choosing a preferable or avoidable antihypertensive medication, or choosing a preferable or avoidable antidiabetic medication, or that said variant gene or materials connected thereto can be utilised as a part of a kit constructed for said uses.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a DNA molecule encoding a variant human α_{2B} -adrenoceptor, said variant α_{2B} -adrenoceptor protein and a method to assess the risk of individuals to develop diabetes or a metabolic syndrome in mammals as well as a method for the targeting treatment for diabetes and selecting subjects for clinical trials.

The word treating shall also be understood to include preventing.

The concept "a deletion of at least 1 glutamate from a glutamic acid repeat element of 12 glutamates" refers to any deletion of 1 to 12 glutamates, amino acids 298–309 (SEQ ID NO: 4), in an acidic stretch of 18 amino acids 294–311 located in the 3rd intracellular loop of the receptor polypeptide irrespective of the specific location of the deletion in said repeat element, or how many glutamates from said repeat element of 12 glutamates are deleted.

The concept "deletion/deletion (D/D) genotype of the human α_{2B} -adrenoceptor", in short "D/D genotype", refers to a genotype of an individual having both α_{2B} -adrenoceptor alleles code for a variant α_{2B} -adrenoceptor with a deletion of at least 1 glutamate from a glutamic acid repeat element of 12 glutamates, amino acids 298–309, in an acidic stretch of 18 amino acids 294–311 (SEQ ID NO: 4), located in the 3rd intracellular loop of the receptor polypeptide. Correspondingly "deletion/insertion (D/I) genotype" refers to a genotype having one of the gene alleles code for an α_{2B} -adrenoceptor with a said deletion and the other without a said deletion, i.e. with a respective insertion, and thus the "insertion/insertion (I/I) genotype" refers to a genotype having both alleles code for an α_{2B} -adrenoceptor without said deletion or deletions.

The term “metabolic syndrome” is defined in the Experimental Section.

A common variant form (SEQ ID NO: 1) of the human α_{2B} -AR gene (SEQ ID NO: 3) has been identified (Heinonen et al. 1999). This variant gene encodes a receptor protein (SEQ ID NO: 2) with a deletion of 3 glutamates, amino acids 307-309, from a glutamic acid (Glu) repeat element of 12 glutamates, amino acids 298–309, in an acidic stretch of 18 amino acids 294–311 (SEQ ID NO: 4), located in the 3rd intracellular loop of the receptor polypeptide. This variant gene (SEQ ID NO: 1) was associated with decreased basal metabolic rate (BMR) in a group of obese Finnish subjects (Heinonen et al. 1999). Of the 166 obese subjects, 47 (28 %) were homozygous for the long 12 glutamate repeat element (Glu¹²/Glu¹²), whereas 90 (54 %) were heterozygous (Glu¹²/Glu⁹) and 29 (17 %) were homozygous for the short form (Glu⁹/Glu⁹).

The results to be presented below show that in a population sample of Finnish middle-aged men subjects homozygous for the short form (Glu⁹/Glu⁹) described above, thus representing a deletion/deletion (D/D) genotype of the α_{2B} -adrenoceptor, have a significantly elevated risk for diabetes. Based on these results and previous publications referred to above it can be postulated that this D/D genotype is related to an impaired capacity to downregulate α_{2B} -adrenoceptor function during sustained receptor activation. Since altered α_{2B} -adrenoceptor function seems to be of relevance in the pathogenesis of diabetes, it could also be of relevance in subjects with the insertion/deletion (I/D) (heterozygous Glu¹²/Glu⁹) and insertion/insertion (I/I) (homozygous Glu¹²/Glu¹²) genotypes when other risk factors for diabetes are present. Further, since this specific deletion of 3 glutamates from said glutamic acid repeat element of 12 glutamates, amino acids 298–309, in said acidic stretch of 18 amino acids 294–311, located in the 3rd intracellular loop of the receptor polypeptide seems to be of relevance in diabetes then also other deletions, i.e. deletions of at least 1 glutamate, from said glutamic acid repeat element of 12 glutamates, amino acids 298–309, could be of relevance in the pathogenesis of diabetes, because the 3rd intracellular loop of the receptor polypeptide said repeat element is located in seems to have an essential role in the down-regulation of the α_{2B} -adrenoceptor. Thus individuals with functional mutations in the α_{2B} -adrenoceptor gene have chronically up-regulated α_{2B} -

adrenoceptors, leading to the vasoconstriction of peripheral arteries and reduced muscle and pancreatic blood flow.

α_{2B} -adrenoceptors mediate contraction of arteries, and genetic polymorphism present in the α_{2B} -adrenoceptor gene renders some subjects more susceptible to α_{2B} -adrenoceptor mediated vasoconstriction of the peripheral blood flow regulating arteries and arterioles and associated clinical disorders such as diabetes or a metabolic syndrome. These subjects will especially benefit from treatment with an α_{2B} -adrenoceptor antagonist, and will be at increased risk for adverse effects if subtype-nonselective α_2 -agonists are administered to them. Therefore, a gene test recognizing subjects with a deletion variant of the α_{2B} -adrenoceptor gene will be useful in diagnostics and patient selection for specific therapeutic procedures and clinical drug testing trials. A gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in assessing an individual's risk to develop diabetes related to the D/D genotype. The test can be used to set a specific subdiagnosis of diabetes, based on its genetic etiology.

Furthermore, a gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in selecting drug therapy for patients with diabetes. Such drugs are e.g. a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptors of the subjects either directly or through central nervous system effects, for example pindolol, propranolol, sotalol, timolol, acebutolol, atenol, betaxolol, bisoprol, esmolol, metoprolol, seliprol, carvedilol, labetalol, clonidine, moxonidine, prazosin, or indapamid, including α -adrenoceptor antagonists (α_{2B} -selective or nonselective).

For instance, as angiotensin II causes an increase of noradrenaline sensitivity, and this effect is at least in part mediated by α -adrenoceptors (Datte et al. 2000), the blood pressure lowering effect of drugs acting through angiotensin II inhibition, such as the angiotensin (AT) receptor blockers, is conceivably enhanced in persons with the D/D genotype of the α_{2B} -adrenoceptor. Such drugs are for example captopril, cinapril, enalapril, imidapril, lisinopril, moexipril, perindopril, ramipril,trandolapril, candesartan, eprosartan, irbesartan, losartan, valsartan or telmisartan.

A gene test recognizing the D/D genotype of the α_{2B} -adrenoceptor is useful in selecting drug therapy for patients who might be at increased risk for adverse effects of α_2 -adrenergic agonists; either it will be possible to avoid the use of α_2 -agonists in such patients, or it will be possible to include a specific α_{2B} -antagonist in their therapeutic regimen.

For instance, a nucleic acid sample from a subject can be used for determining if said subject is a carrier of a variant gene. The determination can be carried out either as a DNA analysis according to well known methods, which include direct DNA sequencing of the normal and variant gene, allele specific amplification using the polymerase chain reaction (PCR) enabling detection of either normal or variant sequence, or by indirect detection of the normal or variant gene by various molecular biology methods including e.g. PCR-single stranded conformation polymorphism (SSCP) or conformational analysis (SSCA) method or denaturing gradient gel electrophoresis (DGGE). Determination of the normal or variant gene can also be done by using a restriction fragment length polymorphism (RFLP) method, which is particularly suitable for genotyping large numbers of samples. Similarly, a test based on gene chip or array technology can be easily developed in analogy with many currently existing such tests for single-nucleotide polymorphisms.

The determination can also be carried out at the level of RNA by analyzing RNA expressed at tissue level using various methods. Allele specific probes can be designed for hybridization. Hybridization can be done e.g. using Northern blot, RNase protection assay or in situ hybridization methods. RNA derived from the normal or variant gene can also be analyzed by converting tissue RNA first to cDNA and thereafter amplifying cDNA by an allele specific PCR method.

The presence of variant α_{2B} -adrenoceptor polypeptides indicating the presence of variant gene can be detected by various methods such as hybridization (Western blot), or other antibody based protein assays.

A kit for detecting a risk of diabetes or a metabolic syndrome preferably contains the various components needed for carrying out the diagnostic method according to the present invention. These components are preferably packaged in separate containers

and/or vials. The kit may also include instructions for carrying out the method. Thus, for example, some or all of the various reagents and other ingredients needed for carrying out the determination, such as buffers, primers, enzymes, control samples or standards etc. can be packaged separately but provided for use in the same box.

- 5 Instructions for carrying out the method can be included inside the box, as a separate insert, or as a label on the box and/or on the separate vials. The kit may also contain the necessary computer software needed to interpret the results obtained with the kit, or for utilizing the results from a gene chip used in the method. Preferably, the kit contains a capturing nucleic acid or an antibody specific to a variant α_{2B} -adrenoceptor nucleic acid
10 or polypeptide, respectively.

The publications and other materials used herein to illuminate the background of the invention, and in particular, to provide additional details with respect to its practice, are incorporated herein by reference.

- 15 The invention will be described in more detail in the experimental section.

EXPERIMENTAL SECTION

Determination of genomic alleles encoding the α_2B -adrenoceptor

DNA fragment analysis of ADRA2B insertion/deletion polymorphism

5 ADRA2B insertion/deletion mutation

The nucleotide sequence of the primer pair for the amplification of the human ADRA2B gene (alpha2B-adrenergic receptor gene) insertion/deletion polymorphism (SEQ ID NO:3) (SEQ ID NO:1) was as follows 5'– GGG TGT TTG TGG GGC ATC TC –3' (SEQ ID NO:5) and 5'– TGG CAC TGC CTG GGG TTC A –3' (SEQ ID NO:6). A fluorescent label has been added to the 5' end of one of the above mentioned PCR primers. Thus, the pcr fragment is detectable in the capillary electrophoresis conducted with ABI Prism 3100 Genetic Analyzer.

The genomic DNA region of the mutation in question can be amplified with PCR with PTC-220 DNA Dyad PCR machine (MJ Research). The PCR reaction was conducted in a 20 μ l volume: the reaction mixture contained 60 ng human genomic DNA (extracted from peripheral blood), 1X PCR Buffer (QIAGEN), 100 μ l of each of the nucleotides (dATP, dCTP, dGTP, dTTP), 0.5 μ M of each of the primers and 1 unit of the DNA polymerase (QIAGEN, Hot Start Taq DNA polymerase). The PCR conditions need to be determined experimentally, and the following standard protocol can be used as a start: first the reaction was hold 7 minutes at 94°C, then the following three steps were repeated for 35 cycles: 45 seconds at 94°C, 45 seconds at 54°C, 1 minute at 72°C, after which the reaction was kept at 72°C for an additional 5 minutes and finally hold at 4°C.

25

The insertion/deletion polymorphism of ADRA2B gene concerns an insertion or an deletion of three glutamic acids in the region of 12 Glu aminoacids in the codons 298-309 (SEQ ID NO:3). Thus depending on the genotype, there is either 9 Glu (deletion) (SEQ ID NO:2) or 12 Glu (insertion) (SEQ ID NO:4) at the ADRA2B protein.

30 Depending on whether the amplified allele had an insertion or a deletion in the studied

locus, the size of the PCR product was 91 bp (insertion allele) or 82 bp (deletion allele). Thus, for homozygotes (insertion/insertion or deletion/deletion) only one size of a fragment was detected either 91 bp or 82 bp, respectively. For heterozygotes both of the above mentioned fragments were detected.

5

Capillary electrophoresis with ABI Prism 3100 Genetic Analyzer

A sample of the ADRA2B insertion/deletion PCR product, 9.00 µl of Hi-Di formamide (Applied Biosystems) and 0.25 µl GeneScan-120 LIZ size standard (Applied Biosystems) were combined in a 96-well 3100 optical microamp plate (Applied Biosystems). The reactions were denatured by placing them at 95°C for 5 minutes and then loaded onto an ABI Prism 3100 Genetic Analyzer (Applied Biosystems). Electrophoresis data was processed and the genotypes were visualized by using the GeneScan Analysis version 3.7 (Applied Biosystems).

Study Population

15 The above referred study population of 1426 Finnish middle-aged men subjects including 304 subjects with a specific deletion/deletion (D/D) genotype of the α_{2B} -adrenoceptor is described in more detail in the following:

Knowing the possible involvement of the investigated acidic region in the desensitization mechanism of the receptor we hypothesized that the observed
20 insertion/deletion allelic variation could be associated with insulin resistance, as indicated by serum insulin concentration, the metabolic syndrome and type 2 diabetes. To test this hypothesis, we carried out a population study in middle-aged Finnish men. The study was carried out as part of the Kuopio Ischemic Heart Disease Risk Factor Study (KIHD), which is an ongoing population-based study designed to investigate risk
25 factors for cardiovascular diseases, type 2 diabetes and related outcomes in men from eastern Finland (Salonen 1988). This area is known for its homogenous population (Sajantila et al. 1996) and high coronary morbidity and mortality rates (Keys 1980).

Body-mass index was computed as the ratio of weight (kg) to the square of height (m). Waist circumference was taken as the average of 2 measurements taken after inspiration
30 and after expiration (mean difference between the two measurements \cong 1.5cm) at the

midpoint between the lowest rib and the iliac crest. Waist-hip ratio was defined as the ratio of waist girth to the circumference of the hips measured at the trochanter major.

Subjects were asked to fast for 12 h before blood sampling. They were also asked to refrain from smoking for 12 h and from consuming alcohol for 3 days before blood
 5 draws. Blood glucose was measured at baseline and 11-year follow-up using a glucose dehydrogenase method after precipitation of proteins by trichloroacetic acid. The serum samples for insulin determination were stored at -80°C. Serum insulin was determined with a Novo Biolabs radioimmunoassay kit (Novo Nordisk, Bagsvaerd, Denmark).

Diabetes was defined as fasting blood glucose of 6.7 mmol/L or more or diagnosed
 10 diabetes with either dietary, oral or insulin treatment. The metabolic syndrome for men according to the WHO definition was modified for epidemiological studies as proposed by the EGIR (Balkau et al. 1999) and defined as: hyperinsulinemia (fasting insulin levels in the top 25% of the non-diabetic population), impaired fasting glycemia or diabetes and the presence of at least two of the following: abdominal obesity,
 15 dyslipidemia (triglycerides ≥ 1.70 or HDL < 0.9 mmol/l), or hypertension (blood pressure $\geq 140/90$ mm Hg or blood pressure medication) (Alberti et al 1998). Insulin resistance was approximated as hyperinsulinemia based on fasting insulin concentrations in the upmost fourth (Balkau et al. 1999). Insulin resistance was also estimated as the bottom fourth of insulin sensitivity as measured by a validated index
 20 (QUICKI) based on fasting insulin and glucose concentrations ($[\log(\text{insulin}) + \log(\text{glucose})]^{-1}$) (Katz et al 2000). Hypertension was defined according to the EGIR recommendations at a lower level than the original WHO definition for consistence with current WHO-ISH and Sixth Joint National Committee recommendations (Balkau et al. 1999, The sixth report 1997, Guidelines Subcommittee 1999). Microalbuminuria
 25 was not included in the definition (Balkau et al. 1999).

Abdominal obesity was defined according to two definitions: 1) according to the original WHO definition - waist-hip ratio > 0.90 or body-mass index ≥ 30 kg/m² (Alberti et al 1998) and 2) modified according to the EGIR recommendation - waist circumference ≥ 94 cm (Balkau et al. 1999).

The metabolic syndrome as defined by the NCEP comprises three or more of the following indications: fasting plasma glucose levels ≥ 6.1 mmol/l (blood glucose levels ≥ 5.6 mmol/l), serum triglycerides ≥ 1.7 mmol/l, serum HDL < 1.0 mmol/l, blood pressure $\geq 135/85$ mmHg, waist girth > 102 cm (Executive Summary 2001). Use of waist girth > 94 cm was suggested for men genetically susceptible to insulin resistance (Executive Summary 2001).

Of all the 1426 subjects, 304 (21%) had the homozygous deletion (D/D) genotype. Of 809 subjects who had no diabetes at the baseline examination and for whom 11-year follow-up information was available, 168 were D/D homozygotes. Of the subjects with D/D genotype, 19 (11.3%) developed diabetes during 11 years of follow-up, whereas of the other subjects, 39 (6.1%) developed diabetes. The odds ratio was 2.0 (95% confidence interval 1.1 to 3.5, $p=0.028$ in 2-sided Fisher's exact test and $p=0.021$ in a logistic model). Other strongest predictors of diabetes during the follow-up were body-mass index (kg/m²) and the waist-to-hip circumference ratio. In a multivariate logistic model including these covariates, the D/D genotype was associated with a 1.9-fold (95% confidence interval 1.1 to 3.5, $p=0.033$) probability (incidence) of diabetes.

The mean fasting baseline serum insulin was 11.1 (mU/L) in 304 D/D homozygotes and 10.5 mU/L in other 1122 genotyped men ($p=0.045$ for difference in Mann-Whitney U test). The D/D genotype of α_{2B} -adrenoceptor gene was also associated with the prevalence and incidence of metabolic syndrome.

The effects of α -adrenoceptor antagonists, β -adrenoceptor antagonists (beta-blockers) and diuretics on serum insulin and lipids and diabetes was analyzed separately in 278 subjects with an antihypertensive medication, of whom 66 were α_{2B} -AR deletion/deletion homozygotes and 212 had other genotypes. Among the non-homozygotes (the wild I allele carriers), the use of α -adrenoceptor antagonists was associated with a lowering of both fasting (6.9 vs. 11.3 mU/L, $p=0.020$) and 2h post glucose load serum insulin concentrations (34.0 vs. 72.3 mU/L, $p<0.001$), while among the D/D homozygotes, α -adrenoceptor antagonists had no effect on either fasting or post-load serum insulin concentration. Among the α_{2B} -AR deletion homozygotes, the use of β -adrenoceptor antagonist therapy was associated with a lowering of serum total

(5.1 vs 5.8 mmol/L, $p=0.010$) and LDL cholesterol (3.6 vs. 4.1 mmol/L, $p=0.053$) concentrations, while among other genotypes, both serum cholesterol and LDL cholesterol tended to be higher among β -blocker users than among non-users. Among the deletion homozygotes, the use of diuretics was associated elevations of serum total
5 (6.0 vs 5.5 mmol/L) and LDL cholesterol (4.4 vs 3.8 mmol/L) and lowering of triglycerides (1.5 vs 2.1 mmol/L, $p=0.005$), whereas among subjects with other genotypes, serum total and LDL cholesterol tended to be lower and triglycerides higher among the diuretic users.

The prevalence of diabetes (fasting blood glucose of 6.0 mmol/L or more or diagnosed
10 diabetes with either dietary, oral or insulin treatment) at the KIHHD 11-year follow-up was in the α_{2B} -AR deletion carriers higher among α -adrenoceptor antagonist users than non-users (25.0% vs 13.2%), while in the non-carriers, the prevalence of diabetes was lower among the users than the non-users (0% vs 12.3%).

Taken together, the known biological properties of the α_{2B} -AR, the homogeneity of the
15 Finnish population, the study design, the relatively large representative study population and the association of diabetes with one trait suggest that the D/D receptor allele is a causal genetic risk factor for diabetes and modifies the effects of α - and β -adrenoceptor modulating drugs and diuretics on glucose and lipid metabolism. The α_{2B} -AR deletion allele homozygosity or carrier status appears to attenuate or abolish the
20 beneficial effect of α -adrenoceptor antagonists on glucose metabolism, to induce cholesterol and LDL cholesterol elevation by diuretics and reverse or attenuate the cholesterol and LDL cholesterol elevating effect of β -blocking agents.

It will be appreciated that the methods of the present invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. It will
25 be apparent for the specialist in the field that other embodiments exist and do not depart from the spirit of the invention. Thus, the described embodiments are illustrative and should not be construed as restrictive.

References

Ahren B. Autonomic regulation of islet hormone secretion--implications for health and disease. *Diabetologia*. 2000; 43: 393-410.

- 5 Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539-53.

- 10 American Diabetes Association. Consensus Development Conference on Insulin Resistance. *Diabetes Care*. 1998; 21: 310-314.

Balkau B, Charles MA. Comment on the provisional report from the WHO consultation. European Group for the Study of Insulin Resistance (EGIR). *Diabet Med* 1999;16:442-3.

- 15 Bengtsson C. Metabolic side effects of diuretics and beta-adrenoceptor blockers. *Acta Pharmacol Toxicol*. 1984; 54 Suppl 1: 67-70.

- 20 Bengtsson C, Blohme G, Lapidus L, Lindquist O, Lundgren H, Nystrom E, Petersen K, Sigurdsson JA. Do antihypertensive drugs precipitate diabetes? *Br Med J (Clin Res Ed)*. 1984; 289: 1495-1497.

- 25 Borchard U. The role of the Sympathetic Nervous System in Cardiovascular Disease. *J Clin Basic Cardiol*. 2001; 4: 175-177.

Calzada BC, Artinano AL. Alpha-adrenoceptor subtypes. *Pharmacol Res* 2001; 44: 195-208.

- 30 Docherty JR. Subtypes of functional α_1 - and α_2 -receptors. *Eur J Pharmacol* 1998; 361: 1-15.

Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). JAMA 2001;285:2486-97.

- 5 Greenberg A. Diuretic complications. Am J Med Sci. 2000; 319: 10-24. Perez-Stable E, Caralis PV. Thiazide-induced disturbances in carbohydrate, lipid, and potassium metabolism. Am Heart J. 1983; 106: 245-251.

- 10 Gress TW, Nieto FJ, Shahar E, Wofford MR, Brancati FL. Hypertension and antihypertensive therapy as risk factors for type 2 diabetes mellitus. Atherosclerosis Risk in Communities Study. N Engl J Med. 2000; 342: 905-912.

- 15 Heinonen P, Koulu M, Pesonen U, Karvonen M, Rissanen A, Laakso M, Valve R, Uusitupa M, Scheinin M. Identification of a three amino acid deletion in the alpha-2B-adrenergic receptor which is associated with reduced basal metabolic rate in obese subjects. J Clin Endocrinol Metab 1999;84:2429-2433.

- 20 Houston MC. Adverse effects of antihypertensive drug therapy on glucose intolerance. Cardiol Clin. 1986; 4: 117-135.

Jousilahti P, Vartiainen E, Tuomilehto J, Pekkanen J, Puska P. Role of known risk factors in explaining the difference in the risk of coronary heart disease between eastern and southwestern Finland. Ann Med 1998; 30: 481-487.

- 25 Julius S, Gudbrandsson T, Jamerson K, Tariq Shahab S, Andersson O. The hemodynamic link between insulin resistance and hypertension. J Hypertens. 1991; 9: 983-986.

Kaplan GA, Salonen JT. Socioeconomic conditions in childhood and ischaemic heart disease during middle age. BMJ 1990; 301: 1121-1123.

Katz A, Nambi SS, Mather K, et al. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 2000;85:2402-10.

- 5 Keys A. Seven Countries: A Multivariate Analysis of Death and Coronary Heart Disease. Cambridge, Mass, Harvard University Press, 1980.

Kvist-Reimer M, Sundler F, Ahren B. Effects of chemical sympathectomy by means of 6-hydroxydopamine on insulin secretion and islet morphology in alloxan-diabetic mice. *Cell Tissue Res*. 2002; 307: 203-209.

10

Laukkanen JA, Kurl S, Lakka TA, et al. Exercise-induced silent myocardial ischemia and coronary morbidity and mortality in middle-aged men. *J Am Coll Cardiol* 2001;38(1):72-79.

- 15 Lithell HO. Insulin resistance and diabetes in the context of treatment of hypertension. *Blood Press*. 1998; Suppl 3: 28-31.

Luna B, Feinglos MN. Drug-induced hyperglycemia. *JAMA*. 2001; 286: 1945-1948.

- 20 MacMillan LB, Lakhani P, Lovinger D, Limbird LE. Alpha 2-adrenergic receptor subtypes: subtle mutation of the alpha 2A-adrenergic receptor in vivo by gene targeting strategies reveals the role of this subtype in multiple physiological settings. *Recent Prog Horm Res*. 1998; 53: 25-42.

- 25 Moan A, Eide IK, Kjeldsen SE. Metabolic and adrenergic characteristics of young men with insulin resistance. *Blood Press Suppl*. 1996; 1: 30-37.

Muntzel MS. Insulin-mediated sympathoexcitation in obesity and type 2 diabetes. *Nephrol Dial Transplant*. 1999; 14: 2282-2285.

Nonogaki K. New insights into sympathetic regulation of glucose and fat metabolism. *Diabetologia*. 2000; 43: 533-549.

5 Papaccio G, Esposito V. Hyperglycemic effects of hydrochlorothiazide and propranolol. A biochemical and ultrastructural study. *Acta Diabetol Lat*. 1987; 24: 325-330.

10 Reaven GM, Lithell H, Landsberg L. Hypertension and associated metabolic abnormalities--the role of insulin resistance and the sympathoadrenal system. *N Engl J Med*. 1996; 334: 374-381.

Sajantila A, Salem AH, Savolainen P, Bauer K, Gierig C, Paabo S: Paternal and maternal DNA lineages reveal a bottleneck in the founding of the Finnish population. *Proc Natl Acad Sci U.S.A.* 1996;93:12035-12039.

15 Salonen JT. Is there a continuing need for longitudinal epidemiologic research? The Kuopio Ischaemic Heart Disease Risk Factor Study. *Ann Clin Res* 1988;20:46-50.

20 Samuelsson O, Hedner T, Berglund G, Persson B, Andersson OK, Wilhelmsen L. Diabetes mellitus in treated hypertension: incidence, predictive factors and the impact of non-selective beta-blockers and thiazide diuretics during 15 years treatment of middle-aged hypertensive men in the Primary Prevention Trial Goteborg, Sweden. *J Hum Hypertens* 1994; 8: 257-263.

25 Sowers JR, Bakris GL. Antihypertensive therapy and the risk of type 2 diabetes mellitus. *N Engl J Med*. 2000; 342: 969-970.

Takahashi T, Nishizawa Y, Emoto M, Kawagishi T, Matsumoto N, Ishimura E, Inaba M, Okuno Y, Shimada H, Morii H. Sympathetic function test of vasoconstrictor changes in foot arteries in diabetic patients. *Diabetes Care*. 1998; 21: 1495-1501.

30 The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *Arch Intern Med* 1997;157:2413-46.

Valensi P, Huard JP, Giroux C, Attali JR. Factors involved in cardiac autonomic neuropathy in diabetic patients. *J Diabetes Complications*. 1997; 11:180-187.

- 5 Verma S, Bhanot S, McNeill JH. Sympathectomy prevents fructose-induced hyperinsulinemia and hypertension. *Eur J Pharmacol*. 1999; 373: R1-4.

- Weinrauch LA, D'Elia JA, Gleason RE, Keough J, Mann D, Kennedy FP. Autonomic function in type I diabetes mellitus complicated by nephropathy. A cross-sectional
10 analysis in the presymptomatic phase. *Am J Hypertens*. 1995; 8: 782-789.

Weston PJ, Gill GV. Is undetected autonomic dysfunction responsible for sudden death in Type 1 diabetes mellitus? The 'dead in bed' syndrome revisited. *Diabet Med*. 1999; 16: 626-631.

- 15 Wilcox CS. Metabolic and adverse effects of diuretics. *Semin Nephrol*. 1999 ; 19: 557-568.

- 1999 World Health Organization-International Society of Hypertension Guidelines for
20 the Management of Hypertension. Guidelines Subcommittee. *J Hypertens* 1999;17:151-83.

CLAIMS

1. A method for diagnosing a susceptibility to diabetes or a metabolic syndrome in a subject by determining the pattern of alleles encoding a variant
5 α_{2B} -adrenoceptor protein, comprising the steps of
 - a) providing a biological sample of the subject to be tested,
 - b) detecting the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in the biological sample, the presence of the variant
10 genotype indicating an increased risk of diabetes or a metabolic syndrome in said subject.
- 15 2. The method according to claim 1, where said variant genotype of the human α_{2B} -adrenoceptor gene is any nucleotide variation in the genomic DNA that affects the structure or function of the mature α_{2B} -adrenoceptor protein.
3. The method according to claim 2, wherein said variant genotype of the human α_{2B} -adrenoceptor is a homozygote deletion form of mutation (D/D).
- 20 4. The method according to claim 1, wherein the detection step is a DNA-assay.
5. The method according to claim 1, wherein the detection step is carried out using a gene or DNA chip, microarray, strip, panel or similar combination of more than one genes, mutations or RNA expressions to be assayed.
- 25 6. The method according to claim 1, wherein the allelic pattern is determined using polymerase chain reaction.
7. The method according to claim 1, wherein the biological sample is any sample that comprises genetic material, preferably a blood sample or buccal swab sample.

8. The method according to claim 1, wherein the detection step is based on a capturing probe, a single strand of cDNA, comprising a nucleotide sequence encoding a variant α_{2B} -adrenoceptor protein or a fragment thereof.
9. The method according to claim 1, wherein the detection step is an analysis
5 of the α_{2B} -adrenoceptor polypeptide or a part thereof.
10. The method according to claim 1, wherein said method is used for determining whether a subject will benefit from a treatment with a drug affecting the norepinephrine sensitivity or sympathetic activity of the subject.
11. The method according to claim 1, wherein said method is used for
10 determining whether a subject will benefit from a treatment with an α - or α_{2B} -adrenoceptor antagonist.
12. The method according to claim 11, wherein said method is used for determining whether the treatment with an α - or α_{2B} -adrenoceptor antagonist will lower or elevate serum insulin concentration.
- 15 13. The method according to claim 1, wherein said method is used for determining whether a subject will be at increased risk of hyperglycemia, hyperinsulinemia, a metabolic syndrome and diabetes if a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptors or a diuretic is administered to the subject.
- 20 14. The method according to claim 1, wherein said method is used for determining whether a subject will response to a treatment with diuretics or a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptors.
- 25 15. The method according to claim 1, wherein said method is used for determining whether the treatment with diuretics and a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenargic receptor will lower or elevate serum and plasma cholesterol, LDL cholesterol and triglyceride concentrations.

16. The method according to any one of claims 13 - 15, wherein said drug is pindolol, propranolol, sotalol, timolol, acebutolol, atenol, betaxolol, bisoprol, esmolol, metoprolol, seliprol, carvedilol, labetalol, clonidine, moxonidine, prazosin, or indapamid.
- 5 17. The method according to claim 1 further comprising a step of selecting a subject of the D/D genotype for clinical drug trials testing antidiabetic and insulin sensitivity improving effects of compounds.
18. The method according to claim 17, wherein the compound to be tested is a
10 subject.
19. The method according to claim 17, wherein the compound to be tested is a drug modulating, inhibiting or activating the vascular alpha- or beta-adrenergic receptors of the subject either directly or through central nervous system.
20. A method for treating a human or animal subject suffering from type 2 diabetes
15 or for treating vascular complications of diabetes, said method comprising a step of reducing the sympathetic tone, lowering blood or tissue norepinephrine or epinephrine concentrations and/or antagonizing α - or α_{2B} -adrenoceptors of the human or animal subject.
- 20 21. The method according to claim 19 or 20, wherein said method comprise a step of administering to a subject a compound reducing the sympathetic tone, lowering blood or tissue norepinephrine or epinephrine concentrations and/or antagonizing α_{2B} -adrenoceptors of the human subject or animal.
- 25 22. The method according to claim 21, wherein said compound is a subtype of a selective or nonselective α_{2B} -adrenoceptor antagonist.
23. The method according to claim 21 or 22, wherein said method is a gene therapy or gene transfer method.

24. The method according to claim 23, wherein the gene to be transferred is a non-variant α_{2B} -adrenoceptor gene or a fragment or derivative thereof.

25. A kit for detecting a risk of diabetes or a metabolic syndrome in a subject, comprising means for determining the pattern of alleles encoding a variant α_{2B} -
5 adrenoceptor in a biological sample from said subject, and optionally computer software to interpret the results of the determination.

26. The kit according to claim 25 for determining the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in said biological sample.

27. The kit according to claim 25 or 26 comprising a capturing nucleic acid
10 probe or an antibody.

(57) ABSTRACT

The invention provides a method and kit for detecting or diagnosing a risk of or predisposition to type 2 diabetes or a metabolic syndrome in a subject, the method comprising the steps of providing a biological sample of the subject to be tested
5 and detecting the presence or absence of a variant genotype of the human α_{2B} -adrenoceptor in the biological sample. The invention also relates to a method for the treatment of type 2 diabetes.

SEQUENCE LISTING

<110> Salonen, Jukka T

Pirskanen, Mia

Tuomainen, Tomi-Pekka

Yunus, Faisel

<120> Method and kit for detecting a risk of diabetes

<130> 39012

<160> 6

<170> PatentIn version 3.1

<210> 1

<211> 1344

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(1344)

<223> ADRA2B variant type sequence

<400> 1

atg gac cac cag gac ccc tac tcc gtg cag gcc aca gcg gcc ata gcg	48
Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala	
1 5 10 15	
gcg gcc atc acc ttc ctc att ctc ttt acc atc ttc ggc aac gct ctg	96
Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu	
20 25 30	
gtc atc ctg gct gtg ttg acc agc cgc tcg ctg cgc gcc cct cag aac	144
Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn	
35 40 45	

ctg ttc ctg gtg tgc ctg gcc gcc gcc gac atc ctg gtg gcc acg ctc Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu 50 55 60	192
atc atc cct ttc tgc ctg gcc aac gag ctg ctg ggc tac tgg tac ttc Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe 65 70 75 80	240
cgg cgc acg tgg tgc gag gtg tac ctg gcg ctc gac gtg ctc ttc tgc Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys 85 90 95	288
acc tgc tcc atc gtg cac ctg tgc gcc atc agc ctg gac cgc tac tgg Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp 100 105 110	336
gcc gtg agc cgc gcg ctg gag tac aac tcc aag cgc acc ccg cgc cgc Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg 115 120 125	384
atc aag tgc atc atc ctc act gtg tgg ctc atc gcc gcc gtc atc tgc Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser 130 135 140	432
ctg ccg ccc ctc atc tac aag ggc gac cag ggc ccc cag ccg cgc ggg Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly 145 150 155 160	480
cgc ccc cag tgc aag ctc aac cag gag gcc tgg tac atc ctg gcc tcc Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser 165 170 175	528
agc atc gga tct ttc ttt gct cct tgc ctc atc atg atc ctt gtc tac Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr 180 185 190	576
ctg cgc atc tac ctg atc gcc aaa cgc agc aac cgc aga ggt ccc agg Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg 195 200 205	624
gcc aag ggg ggg cct ggg cag ggt gag tcc aag cag ccc cga ccc gac Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp 210 215 220	672
cat ggt ggg gct ttg gcc tca gcc aaa ctg cca gcc ctg gcc tct gtg His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val 225 230 235 240	720
gct tct gcc aga gag gtc aac gga cac tgc aag tcc act ggg gag aag Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys 245 250 255	768
gag gag ggg gag acc cct gaa gat act ggg acc cgg gcc ttg cca ccc Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro 260 265 270	816
agt tgg gct gcc ctt ccc aac tca ggc cag ggc cag aag gag ggt gtt Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val 275 280 285	864

tgt ggg gca tct cca gag gat gaa gct gaa gag gag gaa gag gag gag	912
Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu Glu	
290 295 300	
 gag gag tgt gaa ccc cag gca gtg cca gtg tct ccg gcc tca gct tgc	960
Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala Ser Ala Cys	
305 310 315 320	
 agc ccc ccg ctg cag cag cca cag ggc tcc cgg gtg ctg gcc acc cta	1008
Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu Ala Thr Leu	
325 330 335	
 cgt ggc cag gtg ctc ctg ggc agg ggc gtg ggt gct ata ggt ggg cag	1056
Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile Gly Gly Gln	
340 345 350	
 tgg tgg cgt cga agg gcg cac gtg acc cgg gag aag cgc ttc acc ttc	1104
Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg Phe Thr Phe	
355 360 365	
 gtg ctg gct gtg gtc att ggc gtt ttt gtg ctc tgc tgg ttc ccc ttc	1152
Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp Phe Pro Phe	
370 375 380	
 ttc ttc agc tac agc ctg ggc gcc atc tgc ccg aag cac tgc aag gtg	1200
Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His Cys Lys Val	
385 390 395 400	
 ccc cat ggc ctc ttc cag ttc ttc ttc tgg atc ggc tac tgc aac agc	1248
Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr Cys Asn Ser	
405 410 415	
 tca ctg aac cct gtt atc tac acc atc ttc aac cag gac ttc cgc cgt	1296
Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp Phe Arg Arg	
420 425 430	
 gcc ttc cgg agg atc ctg tgc cgc ccg tgg acc cag acg gcc tgg tga	1344
Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr Ala Trp	
435 440 445	

<210> 2

<211> 447

<212> PRT

<213> Homo sapiens

<400> 2

Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala
1 5 10 15

Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu
20 25 30

Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn
 35 40 45

Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu
 50 55 60

Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe
 65 70 75 80

Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys
 85 90 95

Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp
 100 105 110

Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg
 115 120 125

Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser
 130 135 140

Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly
 145 150 155 160

Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser
 165 170 175

Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr
 180 185 190

Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg
 195 200 205

Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp
 210 215 220

His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val
 225 230 235 240

Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys
 245 250 255

Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro
 260 265 270

Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val

275

280

285

Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu
 290 295 300

Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala Ser Ala Cys
 305 310 315 320

Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu Ala Thr Leu
 325 330 335

Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile Gly Gly Gln
 340 345 350

Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg Phe Thr Phe
 355 360 365

Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp Phe Pro Phe
 370 375 380

Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His Cys Lys Val
 385 390 395 400

Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr Cys Asn Ser
 405 410 415

Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp Phe Arg Arg
 420 425 430

Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr Ala Trp
 435 440 445

<210> 3

<211> 1353

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1) .. (1353)

<223> ADRA2B wild type sequence

<400> 3

atg gac cac cag gac ccc tac tcc gtg cag gcc aca gcg gcc ata gcg	48
Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ile Ala	
1 5 10 15	
gcg gcc atc acc ttc ctc att ctc ttt acc atc ttc ggc aac gct ctg	96
Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu	
20 25 30	
gtc atc ctg gct gtg ttg acc agc cgc tgc ctg cgc gcc cct cag aac	144
Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn	
35 40 45	
ctg ttc ctg gtg tgc ctg gcc gcc gcc gac atc ctg gtg gcc acg ctc	192
Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu	
50 55 60	
atc atc cct ttc tgc ctg gcc aac gag ctg ctg ggc tac tgg tac ttc	240
Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe	
65 70 75 80	
cgg cgc acg tgg tgc gag gtg tac ctg gcg ctc gac gtg ctc ttc tgc	288
Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys	
85 90 95	
acc tgc tcc atc gtg cac ctg tgc gcc atc agc ctg gac cgc tac tgg	336
Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp	
100 105 110	
gcc gtg agc cgc gcg ctg gag tac aac tcc aag cgc acc ccg cgc cgc	384
Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg	
115 120 125	
atc aag tgc atc atc ctc act gtg tgg ctc atc gcc gcc gtc atc tgc	432
Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser	
130 135 140	
ctg ccg ccc ctc atc tac aag ggc gac cag ggc ccc cag ccg cgc ggg	480
Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly	
145 150 155 160	
cgc ccc cag tgc aag ctc aac cag gag gcc tgg tac atc ctg gcc tcc	528
Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser	
165 170 175	
agc atc gga tct ttc ttt gct cct tgc ctc atc atg atc ctt gtc tac	576
Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr	
180 185 190	
ctg cgc atc tac ctg atc gcc aaa cgc agc aac cgc aga ggt ccc agg	624
Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg	
195 200 205	
gcc aag ggg ggg cct ggg cag ggt gag tcc aag cag ccc cga ccc gac	672
Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp	
210 215 220	
cat ggt ggg gct ttg gcc tca gcc aaa ctg cca gcc ctg gcc tct gtg	720
His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val	

225	230	235	240	
gct tct gcc aga gag gtc aac gga cac tcg aag tcc act ggg gag aag				768
Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys	245	250	255	
gag gag ggg gag acc cct gaa gat act ggg acc cgg gcc ttg cca ccc				816
Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro	260	265	270	
agt tgg gct gcc ctt ccc aac tca ggc cag ggc cag aag gag ggt gtt				864
Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val	275	280	285	
tgt ggg gca tct cca gag gat gaa gct gaa gag gag gaa gag gag gag				912
Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu Glu	290	295	300	
gag gag gag gaa gag tgt gaa ccc cag gca gtg cca gtg tct ccg gcc				960
Glu Glu Glu Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala	305	310	315	320
tca gct tgc agc ccc ccg ctg cag cag cca cag ggc tcc cgg gtg ctg				1008
Ser Ala Cys Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu	325	330	335	
gcc acc cta cgt ggc cag gtg ctc ctg ggc agg ggc gtg ggt gct ata				1056
Ala Thr Leu Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile	340	345	350	
ggt ggg cag tgg tgg cgt cga agg gcg cac gtg acc cgg gag aag cgc				1104
Gly Gly Gln Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg	355	360	365	
ttc acc ttc gtg ctg gct gtg gtc att ggc gtt ttt gtg ctc tgc tgg				1152
Phe Thr Phe Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp	370	375	380	
ttc ccc ttc ttc ttc agc tac agc ctg ggc gcc atc tgc ccg aag cac				1200
Phe Pro Phe Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His	385	390	395	400
tgc aag gtg ccc cat ggc ctc ttc cag ttc ttc ttc tgg atc ggc tac				1248
Cys Lys Val Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr	405	410	415	
tgc aac agc tca ctg aac cct gtt atc tac acc atc ttc aac cag gac				1296
Cys Asn Ser Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp	420	425	430	
ttc cgc cgt gcc ttc cgg agg atc ctg tgc cgc ccg tgg acc cag acg				1344
Phe Arg Arg Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr	435	440	445	
gcc tgg tga				1353
Ala Trp	450			

<211> 450

<212> PRT

<213> Homo sapiens

<400> 4

Met Asp His Gln Asp Pro Tyr Ser Val Gln Ala Thr Ala Ala Ile Ala
1 5 10 15

Ala Ala Ile Thr Phe Leu Ile Leu Phe Thr Ile Phe Gly Asn Ala Leu
20 25 30

Val Ile Leu Ala Val Leu Thr Ser Arg Ser Leu Arg Ala Pro Gln Asn
35 40 45

Leu Phe Leu Val Ser Leu Ala Ala Ala Asp Ile Leu Val Ala Thr Leu
50 55 60

Ile Ile Pro Phe Ser Leu Ala Asn Glu Leu Leu Gly Tyr Trp Tyr Phe
65 70 75 80

Arg Arg Thr Trp Cys Glu Val Tyr Leu Ala Leu Asp Val Leu Phe Cys
85 90 95

Thr Ser Ser Ile Val His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Trp
100 105 110

Ala Val Ser Arg Ala Leu Glu Tyr Asn Ser Lys Arg Thr Pro Arg Arg
115 120 125

Ile Lys Cys Ile Ile Leu Thr Val Trp Leu Ile Ala Ala Val Ile Ser
130 135 140

Leu Pro Pro Leu Ile Tyr Lys Gly Asp Gln Gly Pro Gln Pro Arg Gly
145 150 155 160

Arg Pro Gln Cys Lys Leu Asn Gln Glu Ala Trp Tyr Ile Leu Ala Ser
165 170 175

Ser Ile Gly Ser Phe Phe Ala Pro Cys Leu Ile Met Ile Leu Val Tyr
180 185 190

Leu Arg Ile Tyr Leu Ile Ala Lys Arg Ser Asn Arg Arg Gly Pro Arg
195 200 205

Ala Lys Gly Gly Pro Gly Gln Gly Glu Ser Lys Gln Pro Arg Pro Asp
 210 215 220

His Gly Gly Ala Leu Ala Ser Ala Lys Leu Pro Ala Leu Ala Ser Val
 225 230 235 240

Ala Ser Ala Arg Glu Val Asn Gly His Ser Lys Ser Thr Gly Glu Lys
 245 250 255

Glu Glu Gly Glu Thr Pro Glu Asp Thr Gly Thr Arg Ala Leu Pro Pro
 260 265 270

Ser Trp Ala Ala Leu Pro Asn Ser Gly Gln Gly Gln Lys Glu Gly Val
 275 280 285

Cys Gly Ala Ser Pro Glu Asp Glu Ala Glu Glu Glu Glu Glu Glu
 290 295 300

Glu Glu Glu Glu Glu Cys Glu Pro Gln Ala Val Pro Val Ser Pro Ala
 305 310 315 320

Ser Ala Cys Ser Pro Pro Leu Gln Gln Pro Gln Gly Ser Arg Val Leu
 325 330 335

Ala Thr Leu Arg Gly Gln Val Leu Leu Gly Arg Gly Val Gly Ala Ile
 340 345 350

Gly Gly Gln Trp Trp Arg Arg Arg Ala His Val Thr Arg Glu Lys Arg
 355 360 365

Phe Thr Phe Val Leu Ala Val Val Ile Gly Val Phe Val Leu Cys Trp
 370 375 380

Phe Pro Phe Phe Phe Ser Tyr Ser Leu Gly Ala Ile Cys Pro Lys His
 385 390 395 400

Cys Lys Val Pro His Gly Leu Phe Gln Phe Phe Phe Trp Ile Gly Tyr
 405 410 415

Cys Asn Ser Ser Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Gln Asp
 420 425 430

Phe Arg Arg Ala Phe Arg Arg Ile Leu Cys Arg Pro Trp Thr Gln Thr
 435 440 445

Ala Trp
450

<210> 5

<211> 20

<212> DNA

<213> Artificial

<220>

<221> misc_feature

<222> (1)..(20)

<223> ADRA2B pcr primer f

<400> 5
gggtgtttgt ggggcatctc

20

<210> 6

<211> 19

<212> DNA

<213> Artificial

<220>

<221> misc_feature

<222> (1)..(19)

<223> ADRA2B pcr primer r

<400> 6
tggcactgcc tggggttca

19